



(11) **EP 0 824 148 A2**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:
18.02.1998 Bulletin 1998/08

(21) Application number: 97113932.4

(22) Date of filing: 13.08.1997

(51) Int. Cl.⁶: **C12N 15/52, C12N 15/60,**
C12N 1/21, C12P 7/62,
C12N 15/74
// (C12N1/21, C12R1:05)

(84) Designated Contracting States:
AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC
NL PT SE

(30) Priority: 14.08.1996 JP 214509/96
25.07.1997 JP 199979/97

(83) Declaration under Rule 28(4) EPC (expert
solution)

(71) Applicant:
The Institute of Physical and Chemical Research
(RIKEN)
Wako-shi, Saitama 351-01 (JP)

(72) Inventors:
• **Toshiaki, Fukui,**
The Inst. of Phys. & Chem. Res.
Wako-shi, Saitama 351-01 (JP)
• **Yoshiharu, Doi,**
The Inst. of Phys. & Chem. Res.
Wako-shi, Saitama 351-01 (JP)

(74) Representative:
Grosse, Rainer, Dipl.-Ing. et al
Gleiss & Grosse
Patentanwaltskanzlei,
Maybachstrasse 6A
70469 Stuttgart (DE)

Remarks:

The applicant has subsequently filed a sequence
listing and declared, that it includes no new matter.

(54) **Polyester synthase gene and process for producing polyester**

(57) The present invention relates to a polyester synthase gene coding for a polypeptide containing the amino acid sequence of SEQ ID NO:2 or a sequence where in said amino acid sequence, one or more amino acids are deleted, replaced or added, said polypeptide bringing about polyester synthase activity; a gene expression cassette comprising the polyester synthase gene and either of open reading frames located upstream and downstream of said gene; a recombinant vector comprising the gene expression cassette; a transformant transformed with the recombinant vector; and a process for producing polyester by culturing the transformant in a medium and recovering polyester from the resulting culture.

EP 0 824 148 A2

Description

Field of the Invention

5 The present invention relates to a polyester synthase gene, a recombinant vector containing the gene, a transformant carrying the recombinant vector, and a process for producing polyester by use of the transformant.

Background of the Invention

10 It is known that a large number of microorganisms biosynthesize poly-3-hydroxybutyrate (P(3HB)) and store it in the form of ultrafine particles as an energy source in the body. P(3HB) extracted from microorganisms is a thermoplastic polymer with a melting temperature of about 180 °C, and because of its excellent biodegradability and biocompatibility it is drawing attention as "green" plastic for preservation of the environment. Further, P(3HB) is "green" plastic which can be synthesized from regenerable carbon resources including sugars and vegetable oils by various microorganisms. However, P(3HB) is a highly crystalline polymer and thus has the problem in physical properties of inferior resistance to impact, so its practical application has never been attempted.

15 Recently, polyester P(3HB-co-3HH) as a random copolymer of 3-hydroxybutyrate (3HB) and 3-hydroxyhexanoate (3HH) and a process for producing the same have been studied and developed, and these are described in e.g. Japanese Patent Laid Open Publication Nos. 93049/1993 and 265065/1995 respectively. In these publications, the P(3HB-co-3HH) copolymer is produced from alkanolic acids or olive oil by fermentation with *Aeromonas caviae* isolated from soil. It is revealed that because the degree of crystallinity of the P(3HB-co-3HH) copolymer produced through fermentation is reduced with an increasing ratio of the 3HH unit in it, so that the copolymer becomes a soft polymeric material excellent in thermostability and formability and can be manufactured into strong yarn or transparent flexible film (Y. Doi, S. Kitamura, H. Abe, *Macromolecules* 28, 4822-4823 (1995)). However, the yield of polyester (content of polyester in dried microorganisms) according to the processes described in Japanese Patent Laid Open Publication Nos. 20 93049/1993 and 265065/1995 is low, and thus there is demand for developments in a process for producing the copolymerized polyester P(3HB-co-3HH).

Summary of the Invention

30 The object of the present invention is to provide a polyester synthase gene, recombinant vectors containing the gene, transformants transformed with the recombinant vectors, and processes for producing polyester by use of the transformants.

35 As a result of their eager research, the present inventors succeeded in producing the polyester in high yield by cloning a polyester synthase gene and deleting one or both of open reading frames located upstream and downstream of said gene to arrive at the completion of the present invention.

40 That is, the present invention is a polyester synthase gene coding for a polypeptide containing the amino acid sequence of SEQ ID NO:2 or a sequence where in said amino acid sequence, one or more amino acids are deleted, replaced or added, said polypeptide bringing about polyester synthase activity. Said gene includes those containing e.g. the nucleotide sequence of SEQ ID NO:1.


45 Further, the present invention is a gene expression cassette comprising said polyester synthase gene and either of open reading frames located upstream and downstream of said gene. In said gene expression cassette, the open reading frame located upstream of the polyester synthase gene includes those (e.g. SEQ ID NO:3) containing DNA coding for the amino acid sequence of SEQ ID NO:4, and the open reading frame located downstream of the polyester synthase gene includes those (e.g. SEQ ID NO:5) containing DNA coding for a polypeptide containing the amino acid sequence of SEQ ID NO:6 or a sequence where in said amino acid sequence, one or more amino acids are deleted, replaced or added, said polypeptide bringing about enoyl-CoA hydratase activity.

50 Even if one or more amino acids in the amino acid sequence of SEQ ID NO:2 have undergone mutations such as deletion, replacement, addition etc., DNA coding for a polypeptide containing said amino acid sequence is also contained in the gene of the present invention insofar as the polypeptide has polyester synthase activity. For example, DNA coding for the amino acid sequence of SEQ ID NO:2 where methionine at the first position is deleted is also contained in the gene of the present invention.

Further, the present invention is recombinant vectors comprising said polyester synthase gene or said gene expression cassette.

55 Further, the present invention is transformants transformed with said recombinant vectors.

Further, the present invention is processes for producing polyester, wherein said transformant is cultured in a medium, and polyester is recovered from the resulting culture. Examples of such polyester are copolymers (e.g. poly(3-hydroxybutyrate-co-3-hydroxyhexanoate) random copolymers) of 3-hydroxyalkanoic acid represented by formula 1:



wherein R represents a hydrogen atom or a C1 to C4 alkyl group.

Brief Description of the Drawing

FIG. 1 shows the structure of the gene of the present invention.

FIG. 2 is a photograph showing the result of SDS-polyacrylamide gel electrophoresis.

Detailed Description of the Invention

Hereinafter, the present invention is described in detail.

(1) Cloning of Polyester synthase gene

The polyester synthase gene of the present invention is separated from a microorganism belonging to the genus Aeromonas.

First, genomic DNA is isolated from a strain having the polyester synthase gene. Such a strain includes e.g. Aeromonas caviae.

Any known methods can be used for preparation of genomic DNA. For example, Aeromonas caviae is cultured in LB medium and then its genomic DNA is prepared by the hexadecyl trimethyl ammonium bromide method (Current Protocols in Molecular Biology, vol. 1, page 2.4.3., John Wiley & Sons Inc., 1994).

The DNA obtained in this manner is partially digested with a suitable restriction enzyme (e.g. Sau3AI, BamHI, BglII etc.) and then the DNA fragments are then dephosphorylated by treatment with alkaline phosphatase. It is ligated into a vector previously cleaved with a restriction enzyme (e.g. BamHI, BglII etc.) to prepare a library.

Phage or plasmid capable of autonomously replicating in host microorganisms is used as the vector. The phage vector includes e.g. EMBL3, M13, λ gt11 etc., and the plasmid vector includes e.g. pBR322, pUC18, and pBluescript II (Stratagene). Vectors capable of autonomously replicating in 2 or more host cells such as E. coli and Bacillus brevis, as well as various shuttle vectors, can also be used. Such vectors are also cleaved with said restriction enzymes so that their fragment can be obtained.

Conventional DNA ligase is used to ligate the resulting DNA fragments into the vector fragment. The DNA fragments and the vector fragment are annealed and then ligated to produce a recombinant vector.

To introduce the recombinant vector into a host microorganism, any known methods can be used. For example, if the host microorganism is E. coli, the calcium method (Lederberg, E.M. et al., J. Bacteriol. 119, 1072 (1974)) and the electroporation method (Current Protocols in Molecular Biology, vol. 1, page 1.8.4 (1994)) can be used. If phage DNA is used, the *in vitro* packaging method (Current Protocols in Molecular Biology, vol. 1., page 5.7.1 (1994)) etc. can be adopted. In the present invention, an *in vitro* packaging kit (Gigapack II, produced by Stratagene etc.) can also be used.

To obtain a DNA fragment containing the polyester synthase gene derived from Aeromonas caviae, a probe is then prepared. The amino acid sequences of some polyester synthase have already been known (Peoples, O.P. and Sinskey, A.J., J. Biol. Chem., 264, 15293 (1989); Huisman, G.W. et al., J. Biol. Chem., 266, 2191 (1991); Pieper, U. et al., FEMS Microbiol. Lett., 96, 73 (1992) etc.). Two conserved regions are selected from these amino acid sequences, and nucleotide sequences coding them are estimated to design oligonucleotides for use as primers. Examples of such oligonucleotides include, but are not limited to, the 2 oligonucleotides 5'-CC(C/G)CC(C/G)TGGATCAA(T/C)AAGT(T/A)(T/C)TA(T/C)ATC-3' (SEQ ID NO:7) and 5'-(G/C)AGCCA (G/C)GC(G/C)GTCCA(A/G)TC(G/C)GGCCACCA-3' (SEQ ID NO:8).

Polymerase chain reaction (PCR) (Molecular Cloning, vol. 2, page 14.2 (1989)) is carried out using these oligonucleotides as primers and the genomic DNA of Aeromonas caviae as a template. The partial fragment of polyester synthase gene is amplified by PCR.

Then, the partially amplified fragment thus obtained is labeled with a suitable reagent and used for colony hybridization of the above genomic DNA library (Current Protocols in Molecular Biology, vol. 1, page 6.0.3 (1994)).

The E. coli is screened by colony hybridization, and a plasmid is recovered from it using the alkaline method (Cur-

rent Protocols in Molecular Biology, vol. 1, page 1.6.1 (1994)), whereby a DNA fragment containing the polyester synthase gene is obtained.

The nucleotide sequence of said DNA fragment can be determined in e.g. an automatic nucleotide sequence analyzer such as 373A DNA sequencer (Applied Biosystems) using a known method such as the Sanger method (Molecular Cloning, vol. 2, page 13.3 (1989)).

The nucleotide sequence of the polyester synthase gene of the present invention is shown in SEQ ID NO:1, and the amino acid sequence encoded by said gene is shown in SEQ ID NO:2, where some amino acids may have undergone mutations such as deletion, replacement, addition etc. insofar as a polypeptide having said amino acid sequence brings about polyester synthase activity. Further, the gene of the present invention encompasses not only the nucleotide sequence coding for the amino acid sequence of SEQ ID NO:2 but also its degenerated isomers which except for degeneracy codons, code for the same polypeptide.

The above mutations such as deletion etc. can be induced by known site-directed mutagenesis (Current Protocols in Molecular Biology, vol. 1, page 8.1.1 (1994)).

After the nucleotide sequence was determined by the means described above, the gene of the present invention can be obtained by chemical synthesis or the PCR technique using genomic DNA as a template, or by hybridization using a DNA fragment having said nucleotide sequence as a probe.

(2) Preparation of Transformant

The transformant of the present invention is obtained by introducing the recombinant vector of the present invention into a host compatible with the expression vector used in constructing said recombinant vector.

The host is not particularly limited insofar as it can express the target gene. Examples are bacteria such as microorganisms belonging to the genus *Alcaligenes*, microorganisms belonging to the genus *Pseudomonas*, microorganisms belonging to the genus *Bacillus*, yeasts such as the genera *Saccharomyces*, *Candida* etc., and animal cells such as COS cells, CHO cells etc.

If bacteria such as microorganisms belonging to the genus *Alcaligenes*, microorganisms belonging to the genus *Pseudomonas* etc. are used as the host, the recombinant DNA of the present invention is preferably constituted such that it contains a promoter, the DNA of the present invention, and a transcription termination sequence so as to be capable of autonomous replication in the host. The expression vector includes pLA2917 (ATCC 37355) containing replication origin RK2 and pJRD215 (ATCC 37533) containing replication origin RSF1010, which are replicated and maintained in a broad range of hosts.

The promoter may be any one if it can be expressed in the host. Examples are promoters derived from *E. coli*, phage etc., such as trp promoter, lac promoter, P_L promoter, P_R promoter and T7 promoter. The method of introducing the recombinant DNA into bacteria includes e.g. a method using calcium ions (Current Protocols in Molecular Biology, vol. 1, page 1.8.1 (1994)) and the electroporation method (Current Protocols in Molecular Biology, vol. 1, page 1.8.4 (1994)).

If yeast is used as the host, expression vectors such as YEp13, YCp50 etc. are used. The promoter includes e.g. gal 1 promoter, gal 10 promoter etc. To method of introducing the recombinant DNA into yeast includes e.g. the electroporation method (Methods. Enzymol., 194, 182-187 (1990)), the spheroplast method (Proc. Natl. Acad. Sci. USA, 84, 1929-1933 (1978)), the lithium acetate method (J. Bacteriol., 153, 163-168 (1983)) etc.

If animal cells are used as the host, expression vectors such as pcDNA1, pcDNA1/Amp (produced by Invitrogen) etc. are used. The method of introducing the recombinant DNA into animal cells includes e.g. the electroporation method, potassium phosphate method etc.

The nucleotide sequence determined as described above contains the polyester synthase gene as well as a plurality of open reading frames (ORFs) upstream and downstream of it. That is, the polyester synthase gene forms an operon with at least 2 ORF's under the control of a single promoter region.

The ORF's which are located respectively upstream and downstream of the polyester synthase gene are referred to hereinafter as "ORF1" and "ORF3".

It is considered that ORF1 is an open reading frame of a gene involved in accumulating polyester in the microorganism or a gene in the polyester biosynthesis system. It was revealed that ORF3 is an open reading frame of a gene coding for enoyl-CoA hydratase (particularly (R)-specific enoyl-CoA hydratase) involved in biosynthesis of polyester.

As shown in FIG. 1, an EcoRI fragment carrying an expression regulatory region (expressed as "-35/-10" in FIG. 1A), the polyester synthase gene, ORF1, and ORF3 was cloned in the present invention (FIG. 1A). This fragment is designated EE32.

Then, a fragment (a gene expression cassette) is prepared by deleting ORF1 and/or ORF3 from EE32, and this cassette is introduced into a host whereby a transformant capable of efficiently producing polyester can be obtained.

In EE32, a restriction enzyme BglII sites are introduced into regions between the expression regulatory region and the translation initiation codon of ORF1 and between the translation termination codon of ORF1 and the translation ini-

tiation codon of the polyester synthase gene, and then ORF1 is deleted from EE32 by treatment with BglII (FIG. 1B). Similarly, a restriction enzyme BamHI sites is introduced into a region between the translation termination codon of the polyester synthase gene and ORF3, and then ORF3 is deleted by treatment with BamHI (FIG. 1C).

To delete both ORF1 and ORF3, EE32 may be subjected to the above operation of deleting ORF1 and ORF3 (FIG. 1D).

The restriction enzyme sites can be introduced by site-directed mutagenesis using synthetic oligonucleotides (Current Protocols in Molecular Biology, vol. 1, page 8.1.1 (1994)).

Each gene expression cassette thus obtained is inserted into said plasmid capable of expression (e.g. pJRD215 (ATCC 37533)) and the resulting recombinant vector is used to transform *Alcaligenes eutrophus* PHB-4 (DSM541) (strain deficient in the ability to synthesize polyester). The method for this transformation includes e.g. the calcium chloride method, rubidium chloride method, low pH method, *in vitro* packaging method, conjugation transfer method etc.

(3) Production of Polyester

The production of polyester is carried out by culturing the transformant of the present invention in a medium, forming and accumulating the polyester of the present invention in the microorganism or in the culture, and recovering the polyester from the cultured microorganism or from the culture.

A conventional method used for culturing the host is also used to culture the transformant of the present invention.

The medium for the transformant prepared from a microorganism belonging to the genus *Alcaligenes* or *Pseudomonas* as the host include a medium containing a carbon source assimilable by the microorganism, in which a nitrogen source, inorganic salts or another organic nutrition source has been limited, for example a medium in which the nutrition source has been limited to 0.01 to 0.1 %.

The carbon source is necessary for growth of the microorganism, and it is simultaneously a starting material of polyester. Examples are hydrocarbons such as glucose, fructose, sucrose, maltose etc. Further, fat and oil related substances having 2 or more carbon atoms can be used as the carbon source. The fat and oil related substances include natural fats and oils, such as corn oil, soybean oil, safflower oil, sunflower oil, olive oil, coconut oil, palm oil, rape oil, fish oil, whale oil, porcine oil and cattle oil, aliphatic acids such as acetic acid, propionic acid, butanoic acid, pentanoic acid, hexoic acid, octanoic acid, decanoic acid, lauric acid, oleic acid, palmitic acid, linolenic acid, linolic acid and myristic acid as well as esters thereof, alcohols such as ethanol, propanol, butanol, pentanol, hexanol, octanol, lauryl alcohol, oleyl alcohol and palmityl alcohol as well as esters thereof.

The nitrogen source includes e.g. ammonia, ammonium salts such as ammonium chloride, ammonium sulfate, ammonium phosphate etc., peptone, meat extract, yeast extract, corn steep liquor etc. The inorganic matter includes e.g. monopotassium phosphate, dipotassium phosphate, magnesium phosphate, magnesium sulfate, sodium chloride etc.

Culture is carried out usually under aerobic conditions with shaking at 25 to 37 °C for more than 24 hours (e.g. 1 to 7 days) after expression is induced. During culture, antibiotics such as ampicillin, kanamycin, antipyrine, tetracycline etc. may be added to the culture. Polyester is accumulated in the microorganism by culturing it, and the polyester is then recovered.

To culture the microorganism transformed with the expression vector using an inducible promoter, its inducer can also be added to the medium. For example, isopropyl- β -D-thiogalactopyranoside (IPTG), indoleacrylic acid (IAA) etc. can be added to the medium.


To culture the transformant from animal cells as the host, use is made of a medium such as RPMI-1640 or DMEM which may be supplemented with fetal bovine serum. Culture is carried out usually in the presence of 5 % CO₂ at 30 to 37°C for 14 to 28 days. During culture, antibiotics such as kanamycin, penicillin etc. may be added to the medium.

In the present invention, purification of polyester can be carried out e.g. as follows:

The transformant is recovered from the culture by centrifugation, then washed with distilled water and dried. Thereafter, the dried transformant is suspended in chloroform and heated to extract polyester from it. The residues are removed by filtration. Methanol is added to this chloroform solution to precipitate polyester. After the supernatant is removed by filtration or centrifugation, the precipitates are dried to give purified polyester.

The resulting polyester is confirmed to be the desired one in a usual manner e.g. by gas chromatography, nuclear magnetic resonance etc.

The gene of the present invention contains the polyester synthase gene isolated from *Aeromonas caviae*. This synthase can synthesize a copolymer (polyester) consisting of a monomer unit 3-hydroxyalkanoic acid represented by formula I:



wherein R represents a hydrogen atom or a C1 to C4 alkyl group. Said copolymer includes e.g. poly(3-hydroxybutyrate-co-3-hydroxyhexanoate) random copolymer (P(3HB-co-3HH)) etc. and the transformant carrying said polyester synthase gene has the ability to produce P(3HB-co-3HH) with very high efficiency.

Conventionally, a process for producing poly-3-hydroxybutyrate (P(3HB)) or poly(3-hydroxybutyrate-co-3-hydroxyvalerate) random copolymer P(3HB-co-3HV) has been studied and developed, but such polyester has the problem in physical properties of inferior resistance to impact because it is a highly crystalline polymer.

Because degree of crystallinity is lowered by introducing 3-hydroxyhexanoate having 6 carbon atoms into a polymer chain, polyester acts as a flexible polymeric material which is also excellent in thermostability and formability, but conventional processes for producing P(3HB-co-3HH) by use of *Aeromonas caviae* (Japanese Patent Laid Open Publication Nos. 93049/1993 and 265065/1995) suffer from a low yield of polyester.

In the present invention, the P(3HB-co-3HH) copolyester can be produced in high yield.

Because the desired polyester can be obtained in a large amount using the above means, it can be used as a biodegradable material of yarn or film, various vessels etc. Further, the gene of the present invention can be used to breed a strain highly producing the P(3HB-co-3HH) copolymer polyester.

Examples

Hereinafter, the present invention is described in more detail with reference to the Examples which however are not intended to limit the scope of the present invention. (Example 1] Cloning of the Polyester synthase Gene from *Aeromonas caviae*

First, a genomic DNA library was prepared from *Aeromonas caviae*.

Aeromonas caviae FA440 was cultured overnight in 100 ml LB medium (1 % yeast extract, 0.5 % trypton, 0.5 % sodium chloride, 0.1 % glucose, pH 7.5) at 30 °C and then genomic DNA was obtained from the microorganism using the hexadecyl trimethyl ammonium bromide method (Current Protocols in Molecular Biology, vol. 1, page 2.4.3 (1994), John Wiley & Sons Inc.).

The resulting genomic DNA was partially digested with restriction enzyme Sau3AI. The vector plasmid used was cosmid vector pLA2917 (ATCC 37355).

This plasmid was cleaved with restriction enzyme BglII and dephosphorylated (Molecular Cloning, vol. 1, page 5.7.2 (1989), Cold Spring Harbor Laboratory) and then ligated into the partially digested genomic DNA fragment by use of DNA ligase.

E. coli S17-1 was transformed with this ligated DNA fragment by the *in vitro* packaging method (Current Protocols in Molecular Biology, vol. 1, page 5.7.2 (1994)) whereby a genomic DNA library from *Aeromonas caviae* was obtained.

To obtain a DNA fragment containing the polyester synthase gene from *Aeromonas caviae*, a probe was then prepared. Two well conserved regions were selected from known amino acid sequences of several polyester synthases, and nucleotide sequences coding for them were estimated, and 2 oligonucleotides 5'-CC(C/G)CC(C/G)TGGATCAA(T/C)AAGT (T/A)(T/C) TA(T/C)ATC-3' (SEQ ID NO:7) and 5'-(G/C)AGCCA(G/C)GC(G/C)GTCCA(A/G)TC(G/C)GGCCACCA-3' (SEQ ID NO:8) were synthesized.

The polyester synthase gene was partially amplified by PCR using these oligonucleotides as primers and the genomic DNA from *Aeromonas caviae* as a template. PCR was carried out using 30 cycles, each consisting of reaction at 94 °C for 30 seconds, 50 °C for 30 seconds, and 72 °C for 60 seconds.

Then, this partially amplified fragment was labeled with digoxigenin using a DIG DNA labeling kit (Boehringer Mannheim) and used as a probe.

Using the probe thus obtained, *E. coli* carrying a plasmid containing the polyester synthase gene was isolated by colony hybridization from the genomic DNA library from *Aeromonas caviae*. By recovering the plasmid from the *E. coli*, a DNA fragment containing the polyester synthase gene was obtained.

The nucleotide sequence of a 3.2 kbp BglII-EcoRI fragment from this fragment was determined by the Sanger method.

As a result, the nucleotide sequence of the 3.2 kb fragment as shown in SEQ ID NOs:9 or 10 was determined.

By further examining homology to this nucleotide sequence, the polyester synthase gene containing the nucleotide sequence (1785 bp) of SEQ ID NO:1 could be identified in this 3.2 kbp nucleotide sequence.

It should be understood that insofar as the protein encoded by the polyester synthase gene of the present invention has the function of gene expression for polyester polymerization, the nucleotide sequence of said gene may have undergone mutations such as deletion, replacement, addition etc.

In a fragment having the nucleotide sequence of SEQ ID NO:9 or 10, a 405 bp gene (ORF3) and a transcription termination region located downstream of the above 1785 bp nucleotide sequence, as well as a 354 bp gene (ORF1) and an expression regulatory region located upstream thereof were identified. The nucleotide sequence of ORF1 is shown in SEQ ID NO:4; the nucleotide sequence of ORF3 in SEQ ID NO:5; and the amino acid sequence encoded by ORF3 in SEQ ID NO: 6.

ORF3 is an open reading frame of a gene coding for enoyl-CoA hydratase involved in biosynthesis of polyester. Insofar as a polypeptide having the amino acid sequence encoded by ORF3 has enoyl-CoA hydratase activity, particularly (R)-specific enoyl-CoA hydratase activity, said amino acid sequence may have undergone mutations such as deletion, replacement and addition of one or more amino acids.

In the nucleotide sequences of SEQ ID NOS:9 and 10, the expression regulatory region is located at the 1- to 383-positions and the transcription termination region at the 3010 to 3187- positions.

[Example 2] Preparation of *Alcaligenes eutrophus* Transformant

The BglII site of the BglII-EcoRI fragment containing this expression regulatory region, ORF1, the polyester synthase gene, ORF3, and the transcriptional termination region was made EcoRI-ended by use of an EcoRI linker whereby a 3.2 kb EcoRI-EcoRI fragment (EE32 fragment) was obtained. This fragment was inserted into plasmid pJRD215 (ATCC 37533) capable of expression in microorganisms belonging to the genus *Alcaligenes*, and the resulting recombinant plasmid was transformed into *Alcaligenes eutrophus* PHB-4 (DSM 541) (strain deficient in the ability to synthesize polyester) by the conjugation transfer method, as follows:

First, the recombinant plasmid was used to transform *E. coli* S17-1 by the calcium chloride method. The recombinant *E. coli* thus obtained and *Alcaligenes eutrophus* PHB-4 were cultured overnight in 1.5 ml LB medium at 30 °C, and the respective cultures, each 0.1 ml, were combined and cultured at 30 °C for 4 hours. This microbial mixture was plated on MBF agar medium (0.9 % disodium phosphate, 0.15 % monopotassium phosphate, 0.05 % ammonium chloride, 0.5 % fructose, 1.5 % agar, 0.3 mg/ml kanamycin) and cultured at 30 °C for 5 days.

Because *Alcaligenes eutrophus* PHB-4 is rendered resistant to kanamycin by transferring the plasmid in the recombinant *E. coli* into it, the colonies grown on the MBF agar medium are a transformant of *Alcaligenes eutrophus*. One colony was isolated from these colonies so that *Alcaligenes eutrophus* AC32 (referred to hereinafter as AC32) was obtained.

AC32 has been deposited as FERM BP-6038 with the National Institute of Bioscience and Human-Technology, Agency of Industrial Science and Technology, Japan.

A restriction enzyme BglII sites were introduced respectively into regions upstream and downstream of the ORF1 gene in the EE32 fragment by site-directed mutagenesis using a synthetic oligonucleotide (Current Protocols in Molecular Biology, vol. 1, page 8.1.1 (1994)), and an ORF1 gene-free fragment was obtained by deleting the BglII-BglII fragment from the EE32 fragment and then inserted into plasmid pJRD215. The resulting recombinant plasmid was used to transform *Alcaligenes eutrophus* PHB-4 by the conjugation transfer method described above. The resulting transformant is referred to hereinafter as AC321.

Similarly, a restriction enzyme BamHI sites were introduced respectively regions upstream and downstream of the ORF3 gene in the EE32 fragment by site-directed mutagenesis, and an ORF3 gene-free fragment was obtained by deleting the BamHI-BamHI fragment from the EE32 fragment and then inserted into plasmid pJRD215. The resulting recombinant plasmid was used to transform *Alcaligenes eutrophus* PHB-4 by the conjugation transfer method described above. The resulting transformant is referred to hereinafter as AC323.

Similarly, a restriction enzyme BglII sites were introduced respectively regions upstream and downstream of the ORF1 gene and a restriction enzyme BamHI sites were introduced respectively regions upstream and downstream of the ORF3 gene in the EE32 fragment, and a gene fragment free of both the ORF1 and ORF3 genes was obtained by deleting the BglII-BglII and BamHI-BamHI fragments from the EE32 fragment and then inserted into plasmid pJRD215. The resulting recombinant plasmid was used to transform *Alcaligenes eutrophus* PHB-4 by the conjugation transfer method described above. The resulting transformant is referred to hereinafter as AC3213.

Further, the polyester synthase gene was amplified by PCR using the EE32 fragment as a template, and the resulting amplification product was inserted into a region between an expression regulatory region and a transcription termination region in a known polyester biosynthesis operon derived from *Alcaligenes eutrophus*. PCR was carried out using 5'-AGTTCGCCGCTCGGGTGTGGGTGAA-3' (SEQ ID NO: 11) and 5'-GGCATATGCGCTCATGCGGCGTCCT-3' (SEQ ID NO: 12) as primers in 30 cycles each consisting of reaction at 94 °C for 30 seconds, 55 °C for 30 seconds and 72 °C for 60 seconds.

This DNA fragment was inserted into plasmid pJRD215, and the resulting plasmid was used to transform *Alcali-*

genes *eutrophus* PHB-4 by the conjugation transfer method described above. The resulting transformant is referred to hereinafter as AC29.

[Example 3] Synthesis of Polyester by *Alcaligenes eutrophus* Transformants

Each of *Alcaligenes eutrophus* H16, PHB-4, AC32, AC321, AC323, AC3213 and AC29 was inoculated into 95 ml MB medium (0.9 % disodium phosphate, 0.15 % monopotassium phosphate, 0.05 % ammonium chloride) containing 1 ml of 1 % sodium octanoate and incubated in a flask at 30 °C. 0.2 g/L kanamycin was contained in the mediums for strains AC32, AC321, AC323, AC3213 and AC29. 12, 24, 36 and 48 hours thereafter, 1 ml of 1 % sodium octanoate was added to each medium (total amount of sodium octanoate added: 0.5 g) and the microorganisms were cultured for 72 hours.

Each of strains H16 and AC3213 was inoculated into the above MB medium to which 1% olive oil, palm oil, corn oil or oleic acid had been added, and each strain was cultured at 30 °C for 72 hours in a flask. 0.2 g/L kanamycin was contained in the mediums for strain AC3213.

Each of strains H16, AC32, AC321, AC323 and AC3213 was inoculated into the above MB medium to which 1% sodium heptanoate had been added, and each strain was cultured at 30 °C in a flask. 0.2 g/L kanamycin was contained in the mediums for strains AC32, AC321, AC323 and AC3213.

While 1 ml of 1% sodium heptanoate was added to each medium (total amount of sodium heptanoate added: 0.5 g) 12, 24, 36 and 48 hours thereafter, the microorganisms were cultured for 72 hours.

The microorganisms were recovered by centrifugation, washed with distilled water and lyophilized, and the weight of the dried microorganisms was determined. 2 ml sulfuric acid/methanol mixture (15 : 85) and 2 ml chloroform were added to 10-30 mg of the dried microorganism, and the sample was sealed and heated at 100 °C for 140 minutes whereby the polyester in the microorganisms was decomposed into methyl ester. 1 ml distilled water was added thereto and stirred vigorously. It was left and separated into 2 layers, and the lower organic layer was removed and analyzed for its components by capillary gas chromatography through a capillary column Neutra BOND-1 (column of 25 m in length, 0.25 mm in inner diameter and 0.4 µm in liquid film thickness, manufactured by GL Science) in Shimadzu GC-14A. The temperature was raised at a rate of 8 °C/min. from an initial temperature of 100 °C. The results are shown in Tables 1, 2 and 3.

Table 1

Synthesis of Polyester Using Octanoic Acid as Carbon Source				
Strain Used <i>A. eutrophus</i>	Weight of Dried Microorganism (g/l)	Content of Polyester in Dried Microorganism (weight-%)	Polyester Comp.	
			3HB	3HH
			(mole-%)	
H16	3.00	86	100	0
PHB-4	0.80	0	-	-
AC32	0.99	33	78	23
AC321	2.85	92	87	13
AC323	2.85	92	88	12
AC3213	3.64	96	85	15
AC29	3.20	94	92	8
3HB: 3-hydroxybutyrate, 3HH: 3-hydroxyhexanoate				

Table 2

Synthesis of Polyester Using Vegetable Oil or Oleic Acid as Carbon Source					
Strain Used <i>A. eutrophus</i>	Carbon Source	Weight of Dried Microorganism (g/l)	Content of Polyester in Dried Microorganism (weight-%)	Polyester Comp.	
				3HB	3HH
				(mole-%)	
H16	olive oil	4.27	79	100	0
	corn oil	3.57	81	100	0
	palm oil	4.13	79	100	0
	oleic acid	4.06	82	100	0
AC3213	olive oil	3.54	76	96	4
	corn oil	3.60	77	95	5
	palm oil	3.58	81	96	4
	oleic acid	2.22	70	96	4

3HB: 3-hydroxybutyrate, 3HH: 3-hydroxyhexanoate

Table 3

Synthesis of Polyester Using Heptanoic Acid as Carbon Source					
Strain Used <i>A. eutrophus</i>	Weight of Dried Microorganism (g/l)	Content of Polyester in Dried Microorganism (weight-%)	Polyester Comp.		
			3HB	3HV	3HHp
			(mole-%)		
H16	2.50	60	50	50	0
AC32	0.77	7	30	67	5
AC321	1.67	55	46	52	2
AC323	1.27	40	48	45	7
AC3213	2.76	67	44	48	8

3HB: 3-hydroxybutyrate, 3HH: 3-hydroxyhexanoate, 3HHp: 3-hydroxyheptanoate

As shown in Table 1, H16 (i.e. wild-type *Alcaligenes eutrophus*) synthesized a poly(3-hydroxybutyrate) homopolymer. This is because 3HH (3-hydroxyhexanoate) having 6 carbon atoms does not serve as a substrate for the polyester synthase possessed by H16. PHB-4 (i.e. the same strain as H16 but deficient in the ability to synthesize polyester) lacks the polyester synthase and thus does not accumulate polyester. AC32 prepared by introducing into PHB-4 the EE32 fragment containing the polyester synthase gene derived from *Aeromonas caviae* accumulated the poly(3-hydroxybutyrate-co-3-hydroxyhexanoate) random copolymer (P(HB-co-3HH)) containing 22 mole-% 3HH (3-hydroxyhexanoate), and this copolymer accounted for 33 % by weight of the dried microorganism.

AC321, AC323 and AC3213 accumulated P(3HB-co-3HH) containing 12 to 15 mole-% 3HH, and the copolymer accounted for 92 to 96 % by weight of the dried microorganisms. As can be seen from these results, the ability of these strains to accumulate polyester was significantly improved by deleting the ORF1 gene and/or ORF3 gene.

P(3HB-co-3HH) was also accumulated in an amount of 94 % by weight of the microorganism even in the case of

AC29 carrying the polyester synthase gene derived from *A. caviae* whose expression regulatory region and transcriptional termination region had been replaced by those derived from *Alcaligenes eutrophus*, indicating that the yield of polyester was significantly improved even using the expression regulatory region and transcriptional termination region of different origin.

When AC3213 producing polyester in the highest yield was cultured using olive oil, corn oil or palm oil as a carbon source, the microorganism accumulated P(3HB-co-3HH) containing 4 to 5 mole-% 3HH, where the copolymer accounted for 76 to 81 % by weight of the microorganism, as shown in Table 2. Even if oleic acid as a fatty acid component contained most abundantly in vegetable oils was used as a carbon source, AC3213 accumulated P(3HB-co-3HH) containing 4 mole-% 3HH, where the copolymer accounted for 70 % by weight of the microorganism. Its corresponding wild strain H16 synthesized only poly(3-hydroxybutyrate) homopolymer under the same conditions.

Alcaligenes eutrophus FA440 is reported to have accumulated 8 % by weight of P(3HB-co-3HH) by use of palmitic acid as a carbon source (Japanese Patent Laid Open Publication No. 265065/1995). On the other hand, the transformant according to the present invention has accumulated 96 % by weight of P(3HB-co-3HH) by use of octanoic acid as a carbon source and 76 to 81 % by weight of P(3HB-co-3HH) by use of extremely cheap vegetable oils as a carbon source, so the comparison therebetween indicates that the method of synthesizing P(3HB-co-3HH) by the transformant used in the present example is an extremely superior method.

When heptanoic acid was used as a carbon source, H16, that is a wild strain of *Alcaligenes eutrophus*, synthesized poly(3-hydroxybutyrate-co-3-hydroxyvalerate) copolymer (P(3HB-co-3HV)). This is because 3HHp (3-hydroxyheptanoate) having 7 carbon atoms does not serve as a substrate for the polyester synthase possessed by H16, AC32, derived from PHB-4 by introduction of the EE32 fragment containing the polyester synthase gene derived from *Aeromonas caviae*, accumulated poly(3-hydroxybutyrate-co-3-hydroxyvalerate-co-3-hydroxyheptanoate) terpolymer (P(3HB-co-3HV-co-3HHp)) containing 5 mole-% 3HHp, where this copolymer accounted for 7 % by weight of the dried microorganism.

Further, each of strains AC321, AC323 and AC3213 accumulated P(3HB-co-3HV-co-3HHp) containing 2 to 8 mole-% 3HHp where the copolymer accounted for 40 to 67 % by weight of the microorganisms, indicating that the yield of polyester was significantly improved by deleting the ORF1 gene and/or ORF3 gene (Table 3).

From these results, it is concluded that copolyesters consisting of 3-hydroxyalkanoic acid with 4 to 7 carbon atoms can be synthesized using the polyester synthase derived from *Aeromonas caviae*.

[Example 4] Identification of Functions of ORF3

The ORF3 gene was amplified by PCR using the EE32 fragment as a template and then inserted into a site downstream of T7 promoter in expression plasmid PET-3a (Novagene). PCR was carried out using 5'-GCCATATGAGCG-CACAATCCCTGGAAGTAG-3' (SEQ ID NO:13) and 5'-CTGGGATCCGCCGGTGCTTAAGGCAGCTTG-3' (SEQ ID NO:14) as primers in 25 cycles each consisting of reaction at 95 °C for 60 seconds and 68 °C for 30 seconds. The resulting plasmid was used to transform *E. coli* BL21 (DE3) (Novagene). The resulting transformant is designated NB3.

NB3 was cultured in LB medium at 30 °C for 4 hours, and isopropyl- β -D-thiogalactopyranoside (IPTG) was added at a final concentration of 0.4 mM to induce expression, and it was further cultured at 30 °C for 2 hours. The microorganism was recovered by centrifugation, disrupted by ultrasonication and centrifuged to give a soluble protein fraction.

As shown in Table 4, high enoyl-CoA hydratase activity was detected in the soluble fraction from the microorganism having the expression plasmid introduced into it.

Table 4
Specific Activity of Enoyl-CoA Hydratase
in Soluble Protein Fraction

	(unit/mg protein)
<i>E. coli</i> BL21/PET-3a	0
<i>E. coli</i> NB3	1700

The enoyl-CoA hydratase activity was determined by measuring a change in absorbance (263 nm) due to double bond hydration, using crotonyl-CoA (Sigma) as substrate (concentration: 0.25 mM). No activity was detected in *E. coli*

into which the control plasmid PET-3a free of the ORF3 gene had been introduced.

Then, the enoyl-CoA hydratase protein was purified. A soluble protein fraction from NB3 was applied to an anion exchange column Q-Sepharose (Pharmacia) and eluted with a gradient of (0 to 1 M) NaCl, and a fraction with enoyl-CoA hydratase activity was collected. SDS-PAGE analysis indicated that the active fraction was homogenous in electrophoresis as shown in FIG. 2. In addition, about 3-fold specific activity could be attained as shown in Table 5.

Table 5

Specific Activity of Enoyl-CoA Hydratase

	(unit/mg protein)
E. coli NB3 soluble protein fraction	1700
anion exchange column elution fraction	5100

The N-terminal amino acid sequence of the enoyl-CoA hydratase protein thus purified was determined. As shown in Table 6, the determined amino acid sequence was the same except for Met in the initiation codon as the amino acid sequence deduced from the nucleotide sequence of the ORF3 gene.

Table 6

Comparison between Amino Acid Sequences

	(unit/mg protein)
N-terminal amino acid sequence of	
purified enoyl-CoA hydratase: SAQSLEVGQKARLSKRFGAA (SEQ ID NO:15)	
amino acid sequence deduced from	
ORF3 nucleotide sequence: MSAQSLEVGQKARLSKRFGAA (SEQ ID NO:16)	

From this, it could be confirmed that ORF3 codes for enoyl-CoA hydratase. It is considered that Met was released by post-translational modification.

Further, the stereospecificity of enoyl-CoA hydratase encoded by ORF3 was examined as follows:

By adding (S)-3-hydroxybutyryl-CoA dehydrogenase (Sigma) (final concentration: 0.2 unit/ml) and oxidized nicotinamide adenine dinucleotide (NAD⁺) (final concentration: 0.5 mM) to a reaction solution for activity measurement, (S)-3-hydroxybutyryl-CoA formed is oxidized to acetoacetyl-CoA by the action of the dehydrogenase if the enoyl-CoA hydratase is specific to the (S)-isomer. During this reaction, NAD⁺ is reduced to form NADH resulting in the generation of a specific absorption at 340 nm. If enoyl-CoA hydratase is specific to the (R)-isomer, NADH is not formed.

As shown in Table 7, the change in absorbance at 340 nm when enoyl-CoA hydratase encoded by ORF3 was used, was the same as in the case where enoyl-CoA hydratase was absent, but if commercially available (S)-specific enoyl-CoA hydratase (Sigma) was used, a change in absorbance due to formation of NADH was observed.

Table 7

Change in Absorbance at 340 nm after 1 Min.	
no addition of enoyl-CoA hydratase	0.045
ORF3-derived enoyl-CoA hydratase	0.047
(S)-isomer specific enoyl-CoA hydratase (Sigma)	0.146

From this result, it was made evident that the purified enoyl-CoA hydratase is specific to the (R)-isomer. Thus, it was found that ORF3 codes for (R)-isomer specific enoyl-CoA hydratase.

According to the present invention, there are provided a polyester synthase, a recombinant vector carrying the gene, a transformant carrying the recombinant vector and a process for producing polyester by use of the transformant.

The present invention is extremely useful in that the present gene codes for a polyester synthase capable of synthesizing polyester as a copolymer consisting of a monomer unit represented by 3-hydroxyalkanoic acid having 4 to 7 carbon atoms, and that the present process can synthesize a biodegradable plastic P(3HB-co-3HH) very efficiently which is excellent in thermostability and formability.

EP 0 824 148 A2

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

(A) NAME: THE INSTITUTE OF PHYSICAL AND CHEMICAL RESEARCH
 (B) STREET: Hirosawa 2-1
 (C) CITY: Wako-shi
 (D) STATE: Saitama
 (E) COUNTRY: Japan
 (F) POSTAL CODE (ZIP): 351-01
 (G) TELEPHONE: 81-48-467-9263
 (H) TELEFAX: 81-48-462-4609

(ii) TITLE OF INVENTION: POLYESTER SYNTHASE GENE AND PROCESS FOR PRODUCING POLYESTER

(iii) NUMBER OF SEQUENCES: 16

(iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
 (B) COMPUTER: IBM PC compatible
 (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 (D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

(v) CURRENT APPLICATION DATA:

APPLICATION NUMBER: 97113932.4

(vi) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: JP 214509/1996
 (B) FILING DATE: 14-AUG-1996

(vi) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: JP 199979/1997
 (B) FILING DATE: 25-JUL-1997

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1785 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(ix) FEATURE:

(A) NAME/KEY: CDS
 (B) LOCATION: 1..1782

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

ATG	AGC	CAA	CCA	TCT	TAT	GGC	CCG	CTG	TTC	GAG	GCC	CTG	GCC	CAC	TAC	48
Met	Ser	Gln	Pro	Ser	Tyr	Gly	Pro	Leu	Phe	Glu	Ala	Leu	Ala	His	Tyr	
1				5				10				15				
AAT	GAC	AAG	CTG	CTG	GCC	ATG	GCC	AAG	GCC	CAG	ACA	GAG	CGC	ACC	GCC	96
Asn	Asp	Lys	Leu	Leu	Ala	Met	Ala	Lys	Ala	Gln	Thr	Glu	Arg	Thr	Ala	
		20						25				30				
CAG	GCG	CTG	CTG	CAG	ACC	AAT	CTG	GAC	GAT	CTG	GGC	CAG	GTG	CTG	GAG	144
Gln	Ala	Leu	Leu	Gln	Thr	Asn	Leu	Asp	Asp	Leu	Gly	Gln	Val	Leu	Glu	

EP 0 824 148 A2

55

EP 0 824 148 A2

5 Gln Leu Ala Val Ser Phe Ser Leu Leu Arg Glu Asn Ser Leu Tyr Trp
385 390 395 400
AAC TAC TAC ATC GAC AGC TAC CTC AAG GGT CAG AGC CCG GTG GCC TTC 1248
Asn Tyr Tyr Ile Asp Ser Tyr Leu Lys Gly Gln Ser Pro Val Ala Phe
405 410 415
GAT CTG CTG CAC TGG AAC AGC GAC AGC ACC AAT GTG GCG GGC AAG ACC 1296
Asp Leu Leu His Trp Asn Ser Asp Ser Thr Asn Val Ala Gly Lys Thr
420 425 430
CAC AAC AGC CTG CTG CGC CGT CTC TAC CTG GAG AAC CAG CTG GTG AAG 1344
His Asn Ser Leu Leu Arg Arg Leu Tyr Leu Glu Asn Gln Leu Val Lys
435 440 445
GGG GAG CTC AAG ATC CGC AAC ACC CGC ATC GAT CTC GGC AAG GTG AAG 1392
Gly Glu Leu Lys Ile Arg Asn Thr Arg Ile Asp Leu Gly Lys Val Lys
450 455 460
ACC CCT GTG CTG CTG GTG TCG GCG GTG GAC GAT CAC ATC GCC CTC TGG 1440
Thr Pro Val Leu Leu Val Ser Ala Val Asp Asp His Ile Ala Leu Trp
465 470 475 480
CAG GGC ACC TGG CAG GGC ATG AAG CTG TTT GGC GGG GAG CAG CGC TTC 1488
Gln Gly Thr Trp Gln Gly Met Lys Leu Phe Gly Gly Glu Gln Arg Phe
485 490 495
CTC CTG GCG GAG TCC GGC CAC ATC GCC GGC ATC ATC AAC CCG CCG GCC 1536
Leu Leu Ala Glu Ser Gly His Ile Ala Gly Ile Ile Asn Pro Pro Ala
500 505 510
GCC AAC AAG TAC GGC TTC TGG CAC AAC GGG GCC GAG GCC GAG AGC CCG 1584
Ala Asn Lys Tyr Gly Phe Trp His Asn Gly Ala Glu Ala Glu Ser Pro
515 520 525
GAG AGC TGG CTG GCA GGG GCG ACG CAC CAG GGC GGC TCC TGG TGG CCC 1632
Glu Ser Trp Leu Ala Gly Ala Thr His Gln Gly Gly Ser Trp Trp Pro
530 535 540
GAG ATG ATG GGC TTT ATC CAG AAC CGT GAC GAA GGG TCA GAG CCC GTC 1680
Glu Met Met Gly Phe Ile Gln Asn Arg Asp Glu Gly Ser Glu Pro Val
545 550 555 560
CCC GCG CGG GTC CCG GAG GAA GGG CTG GCC CCC GCC CCC GGC CAC TAT 1728
Pro Ala Arg Val Pro Glu Glu Gly Leu Ala Pro Ala Pro Gly His Tyr
565 570 575
GTC AAG GTG CCG CTC AAC CCC GTG TTT GCC TGC CCA ACA GAG GAG GAC 1776
Val Lys Val Arg Leu Asn Pro Val Phe Ala Cys Pro Thr Glu Glu Asp
580 585 590
GCC GCA TGA 1785
Ala Ala

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 594 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

45 Met Ser Gln Pro Ser Tyr Gly Pro Leu Phe Glu Ala Leu Ala His Tyr
1 5 10 15
Asn Asp Lys Leu Leu Ala Met Ala Lys Ala Gln Thr Glu Arg Thr Ala
20 25 30
Gln Ala Leu Leu Gln Thr Asn Leu Asp Asp Leu Gly Gln Val Leu Glu
35 40 45
60 Gln Gly Ser Gln Gln Pro Trp Gln Leu Ile Gln Ala Gln Met Asn Trp
50 55 60
Trp Gln Asp Gln Leu Lys Leu Met Gln His Thr Leu Leu Lys Ser Ala
65 70 75 80

EP 0 824 148 A2

Gly Gln Pro Ser Glu Pro Val Ile Thr Pro Glu Arg Ser Asp Arg Arg
 85 90 95
 Phe Lys Ala Glu Ala Trp Ser Glu Gln Pro Ile Tyr Asp Tyr Leu Lys
 100 105 110
 5 Gln Ser Tyr Leu Leu Thr Ala Arg His Leu Leu Ala Ser Val Asp Ala
 115 120 125
 Leu Glu Gly Val Pro Gln Lys Ser Arg Glu Arg Leu Arg Phe Phe Thr
 130 135 140
 Arg Gln Tyr Val Asn Ala Met Ala Pro Ser Asn Phe Leu Ala Thr Asn
 145 150 155 160
 10 Pro Glu Leu Leu Lys Leu Thr Leu Glu Ser Asp Gly Gln Asn Leu Val
 165 170 175
 Arg Gly Leu Ala Leu Leu Ala Glu Asp Leu Glu Arg Ser Ala Asp Gln
 180 185 190
 Leu Asn Ile Arg Leu Thr Asp Glu Ser Ala Phe Glu Leu Gly Arg Asp
 195 200 205
 15 Leu Ala Leu Thr Pro Gly Arg Val Val Gln Arg Thr Glu Leu Tyr Glu
 210 215 220
 Leu Ile Gln Tyr Ser Pro Thr Thr Glu Thr Val Gly Lys Thr Pro Val
 225 230 235 240
 Leu Ile Val Pro Pro Phe Ile Asn Lys Tyr Tyr Ile Met Asp Met Arg
 245 250 255
 20 Pro Gln Asn Ser Leu Val Ala Trp Leu Val Ala Gln Gly Gln Thr Val
 260 265 270
 Phe Met Ile Ser Trp Arg Asn Pro Gly Val Ala Gln Ala Gln Ile Asp
 275 280 285
 Leu Asp Asp Tyr Val Val Asp Gly Val Ile Ala Ala Leu Asp Gly Val
 290 295 300
 25 Glu Ala Ala Thr Gly Glu Arg Glu Val His Gly Ile Gly Tyr Cys Ile
 305 310 315 320
 Gly Gly Thr Ala Leu Ser Leu Ala Met Gly Trp Leu Ala Ala Arg Arg
 325 330 335
 Gln Lys Gln Arg Val Arg Thr Ala Thr Leu Phe Thr Thr Leu Leu Asp
 340 345 350
 Phe Ser Gln Pro Gly Glu Leu Gly Ile Phe Ile His Glu Pro Ile Ile
 355 360 365
 30 Ala Ala Leu Glu Ala Gln Asn Glu Ala Lys Gly Ile Met Asp Gly Arg
 370 375 380
 Gln Leu Ala Val Ser Phe Ser Leu Leu Arg Glu Asn Ser Leu Tyr Trp
 385 390 395 400
 Asn Tyr Tyr Ile Asp Ser Tyr Leu Lys Gly Gln Ser Pro Val Ala Phe
 405 410 415
 35 Asp Leu Leu His Trp Asn Ser Asp Ser Thr Asn Val Ala Gly Lys Thr
 420 425 430
 His Asn Ser Leu Leu Arg Arg Leu Tyr Leu Glu Asn Gln Leu Val Lys
 435 440 445
 Gly Glu Leu Lys Ile Arg Asn Thr Arg Ile Asp Leu Gly Lys Val Lys
 450 455 460
 40 Thr Pro Val Leu Leu Val Ser Ala Val Asp Asp His Ile Ala Leu Trp
 465 470 475 480
 Gln Gly Thr Trp Gln Gly Met Lys Leu Phe Gly Gly Glu Gln Arg Phe
 485 490 495
 Leu Leu Ala Glu Ser Gly His Ile Ala Gly Ile Ile Asn Pro Pro Ala
 500 505 510
 45 Ala Asn Lys Tyr Gly Phe Trp His Asn Gly Ala Glu Ala Glu Ser Pro
 515 520 525
 Glu Ser Trp Leu Ala Gly Ala Thr His Gln Gly Gly Ser Trp Trp Pro
 530 535 540
 Glu Met Met Gly Phe Ile Gln Asn Arg Asp Glu Gly Ser Glu Pro Val
 545 550 555 560
 50 Pro Ala Arg Val Pro Glu Glu Gly Leu Ala Pro Ala Pro Gly His Tyr
 565 570 575
 Val Lys Val Arg Leu Asn Pro Val Phe Ala Cys Pro Thr Glu Glu Asp
 580 585 590
 Ala Ala

55

EP 0 824 148 A2

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 354 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..351

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

```

ATG ATG AAT ATG GAC GTG ATC AAG AGC TTT ACC GAG CAG ATG CAA GGC   48
Met Met Asn Met Asp Val Ile Lys Ser Phe Thr Glu Gln Met Gln Gly
1      5      10      15
TTC GCC GCC CCC CTC ACC CGC TAC AAC CAG CTG CTG GCC AGC AAC ATC   96
Phe Ala Ala Pro Leu Thr Arg Tyr Asn Gln Leu Leu Ala Ser Asn Ile
20     20     25     30
GAA CAG CTG ACC CGG TTG CAG CTG GCC TCC GCC AAC GCC TAC GCC GAA  144
Glu Gln Leu Thr Arg Leu Gln Leu Ala Ser Ala Asn Ala Tyr Ala Glu
35     40     45
CTG GGC CTC AAC CAG TTG CAG GCC GTG AGC AAG GTG CAG GAC ACC CAG   192
Leu Gly Leu Asn Gln Leu Gln Ala Val Ser Lys Val Gln Asp Thr Gln
50     55     60
AGC CTG GCG GCC CTG GGC ACA GTG CAA CTG GAG ACC GCC AGC CAG CTC   240
Ser Leu Ala Ala Leu Gly Thr Val Gln Leu Glu Thr Ala Ser Gln Leu
65     70     75     80
TCC CGC CAG ATG CTG GAT GAC ATC CAG AAG CTG AGC GCC CTC GGC CAG   288
Ser Arg Gln Met Leu Asp Asp Ile Gln Lys Leu Ser Ala Leu Gly Gln
85     90     95
CAG TTC AAG GAA GAG CTG GAT GTC CTG ACC GCA GAC GGC ATC AAG AAA   336
Gln Phe Lys Glu Leu Asp Val Leu Thr Ala Asp Gly Ile Lys Lys
100    105    110
AGC ACG GGC AAG GCC TGA                                     354
Ser Thr Gly Lys Ala
115

```

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 117 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

```

Met Met Asn Met Asp Val Ile Lys Ser Phe Thr Glu Gln Met Gln Gly
1      5      10      15
Phe Ala Ala Pro Leu Thr Arg Tyr Asn Gln Leu Leu Ala Ser Asn Ile
20     25     30
Glu Gln Leu Thr Arg Leu Gln Leu Ala Ser Ala Asn Ala Tyr Ala Glu
35     40     45
Leu Gly Leu Asn Gln Leu Gln Ala Val Ser Lys Val Gln Asp Thr Gln
50     55     60
Ser Leu Ala Ala Leu Gly Thr Val Gln Leu Glu Thr Ala Ser Gln Leu

```

EP 0 824 148 A2

65 70 75 80
 Ser Arg Gln Met Leu Asp Asp Ile Gln Lys Leu Ser Ala Leu Gly Gln
 85 90 95
 Gln Phe Lys Glu Leu Asp Val Leu Thr Ala Asp Gly Ile Lys Lys
 100 105 110
 Ser Thr Gly Lys Ala
 115

2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 405 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..402

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

ATG	AGC	GCA	CAA	TCC	CTG	GAA	GTA	GGC	CAG	AAG	GCC	CGT	CTC	AGC	AAG	48
Met	Ser	Ala	Gln	Ser	Leu	Glu	Val	Gly	Gln	Lys	Ala	Arg	Leu	Ser	Lys	
1				5				10					15			
CGG	TTC	GGG	GCG	GCG	GAG	GTA	GCC	GCC	TTC	GCC	GCG	CTC	TCG	GAG	GAC	96
Arg	Phe	Gly	Ala	Ala	Glu	Val	Ala	Ala	Phe	Ala	Ala	Leu	Ser	Glu	Asp	
			20				25					30				
TTC	AAC	CCC	CTG	CAC	CTG	GAC	CCG	GCC	TTC	GCC	GCC	ACC	ACG	GCG	TTC	144
Phe	Asn	Pro	Leu	His	Leu	Asp	Pro	Ala	Phe	Ala	Ala	Thr	Thr	Ala	Phe	
			35			40						45				
GAG	CGG	CCC	ATA	GTC	CAC	GGC	ATG	CTG	CTC	GCC	AGC	CTC	TTC	TCC	GGG	192
Glu	Arg	Pro	Ile	Val	His	Gly	Met	Leu	Leu	Ala	Ser	Leu	Phe	Ser	Gly	
			50			55					60					
CTG	CTG	GGC	CAG	CAG	TTG	CCG	GGC	AAG	GGG	AGC	ATC	TAT	CTG	GGT	CAA	240
Leu	Leu	Gly	Gln	Gln	Leu	Pro	Gly	Lys	Gly	Ser	Ile	Tyr	Leu	Gly	Gln	
65				70				75					80			
AGC	CTC	AGC	TTC	AAG	CTG	CCG	GTC	TTT	GTC	GGG	GAC	GAG	GTG	ACG	GCC	288
Ser	Leu	Ser	Phe	Lys	Leu	Pro	Val	Phe	Val	Gly	Asp	Glu	Val	Thr	Ala	
			85					90				95				
GAG	GTG	GAG	GTG	ACC	GCC	CTT	CGC	GAG	GAC	AAG	CCC	ATC	GCC	ACC	CTG	336
Glu	Val	Glu	Val	Thr	Ala	Leu	Arg	Glu	Asp	Lys	Pro	Ile	Ala	Thr	Leu	
			100				105					110				
ACC	ACC	CGC	ATC	TTC	ACC	CAA	GGC	GGC	GCC	CTC	GCC	GTG	ACG	GGG	GAA	384
Thr	Thr	Arg	Ile	Phe	Thr	Gln	Gly	Gly	Ala	Leu	Ala	Val	Thr	Gly	Glu	
			115			120						125				
GCC	GTG	GTC	AAG	CTG	CCT	TAA										405
Ala	Val	Val	Lys	Leu	Pro											
			130													

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 134 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS:
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

5 Met Ser Ala Gln Ser Leu Glu Val Gly Gln Lys Ala Arg Leu Ser Lys
 1 5 10 15
 Arg Phe Gly Ala Ala Glu Val Ala Ala Phe Ala Ala Leu Ser Glu Asp
 20 25 30
 Phe Asn Pro Leu His Leu Asp Pro Ala Phe Ala Ala Thr Thr Ala Phe
 35 40 45
 Glu Arg Pro Ile Val His Gly Met Leu Leu Ala Ser Leu Phe Ser Gly
 50 55 60
 10 Leu Leu Gly Gln Gln Leu Pro Gly Lys Gly Ser Ile Tyr Leu Gly Gln
 65 70 75 80
 Ser Leu Ser Phe Lys Leu Pro Val Phe Val Gly Asp Glu Val Thr Ala
 85 90 95
 Glu Val Glu Val Thr Ala Leu Arg Glu Asp Lys Pro Ile Ala Thr Leu
 100 105 110
 15 Thr Thr Arg Ile Phe Thr Gln Gly Gly Ala Leu Ala Val Thr Gly Glu
 115 120 125
 Ala Val Val Lys Leu Pro
 130

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "synthetic DNA"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

CCSCCSTGGA TCAAYAAGTW YTAYATC

(2) INFORMATION FOR SEQ ID NO: 8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "synthetic DNA"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

SAGCCASGCS GTCCARTCSG GCCACCA

(2) INFORMATION FOR SEQ ID NO: 9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 3187 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(ix) FEATURE:

EP 0 824 148 A2

(A) NAME/KEY: CDS
(B) LOCATION: 384..734

(ix) FEATURE:

(A) NAME/KEY: CDS
(B) LOCATION: 830..2611

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

```

AGATCTGGAC CGGGGTGCTG GCCTGGGCCA CGCCGGCGAG GGCCAGCGCG GAGCAACCGA 60
GCAGCAGGGC GAGAGGTTTC ATCGGGATTTC CTTGGCAGTC TGAATGACGT GCCAGCCTAT 120
CAGCGCGGCG CCGGTGCGGC GAGGGCGCGC CGGACCCAGT GCGTCACCTC TCGTCTGATC 180
CGCCTCCCTC GACGGGCGTC GCTGACAAAA AAATTCAAAC AGAAATTAAC ATTTATGTCA 240
TTTACACCAA ACCGCATTG GTTGCAGAAT GCTCAACGT GTGTTTGAAC AGAGCAAGCA 300
ACACGTAAAC AGGGATGACA TCCAGTACCC GTAAGAAGGG CCGATTGGCC CACAACAACA 360
CTGTTCTGCC GAACTGGAGA CCG ATG ATG AAT ATG GAC GTG ATC AAG AGC 410
Met Met Asn Met Asp Val Ile Lys Ser
1 5
TTT ACC GAG CAG ATG CAA GGC TTC GCC GCC CCC CTC ACC CGC TAC AAC 458
Phe Thr Glu Gln Met Gln Gly Phe Ala Ala Pro Leu Thr Arg Tyr Asn
10 15 20 25
CAG CTG CTG GCC AGC AAC ATC GAA CAG CTG ACC CGG TTG CAG CTG GCC 506
Gln Leu Leu Ala Ser Asn Ile Glu Gln Leu Thr Arg Leu Gln Leu Ala
20 30 35 40
TCC GCC AAC GCC TAC GCC GAA CTG GGC CTC AAC CAG TTG CAG GCC GTG 554
Ser Ala Asn Ala Tyr Ala Glu Leu Gly Leu Asn Gln Leu Gln Ala Val
45 50 55
AGC AAG GTG CAG GAC ACC CAG AGC CTG GCG GCC CTG GGC ACA GTG CAA 602
Ser Lys Val Gln Asp Thr Gln Ser Leu Ala Ala Leu Gly Thr Val Gln
60 65 70
CTG GAG ACC GCC AGC CAG CTC TCC CGC CAG ATG CTG GAT GAC ATC CAG 650
Leu Glu Thr Ala Ser Gln Leu Ser Arg Gln Met Leu Asp Asp Ile Gln
75 80 85
AAG CTG AGC GCC CTC GGC CAG CAG TTC AAG GAA GAG CTG GAT GTC CTG 698
Lys Leu Ser Ala Leu Gly Gln Gln Phe Lys Glu Glu Leu Asp Val Leu
90 95 100 105
ACC GCA GAC GGC ATC AAG AAA AGC ACG GGC AAG GCC TGATAACCCC 744
Thr Ala Asp Gly Ile Lys Lys Ser Thr Gly Lys Ala
110 115
TGGTGCCCG TCGGGCAGC CACATCTCCC CATGACTCGA CGCTACGGGC TAGTTCCCGC 804
CTCGGGTGTG GGTGAAGGAG AGCAC ATG AGC CAA CCA TCT TAT GGC CCG CTG 856
Met Ser Gln Pro Ser Tyr Gly Pro Leu
1 5
TTC GAG GCC CTG GCC CAC TAC AAT GAC AAG CTG CTG GCC ATG GCC AAG 904
Phe Glu Ala Leu Ala His Tyr Asn Asp Lys Leu Leu Ala Met Ala Lys
10 15 20 25
GCC CAG ACA GAG CGC ACC GCC CAG GCG CTG CTG CAG ACC AAT CTG GAC 952
Ala Gln Thr Glu Arg Thr Ala Gln Ala Leu Leu Gln Thr Asn Leu Asp
30 35 40
GAT CTG GGC CAG GTG CTG GAG CAG GGC AGC CAG CAA CCC TGG CAG CTG 1000
Asp Leu Gly Gln Val Leu Glu Gln Gly Ser Gln Gln Pro Trp Gln Leu
45 50 55
ATC CAG GCC CAG ATG AAC TGG TGG CAG GAT CAG CTC AAG CTG ATG CAG 1048
Ile Gln Ala Gln Met Asn Trp Trp Gln Asp Gln Leu Lys Leu Met Gln
60 65 70
CAC ACC CTG CTC AAA AGC GCA GGC CAG CCG AGC GAG CCG GTG ATC ACC 1096
His Thr Leu Leu Lys Ser Ala Gly Gln Pro Ser Glu Pro Val Ile Thr
75 80 85
CCG GAG CGC AGC GAT CGC CGC TTC AAG GCC GAG GCC TGG AGC GAA CAA 1144
Pro Glu Arg Ser Asp Arg Arg Phe Lys Ala Glu Ala Trp Ser Glu Gln
90 95 100 105
CCC ATC TAT GAC TAC CTC AAG CAG TCC TAC CTG CTC ACC GCC AGG CAC 1192
Pro Ile Tyr Asp Tyr Leu Lys Gln Ser Tyr Leu Leu Thr Ala Arg His
110 115 120
CTG CTG GCC TCG GTG GAT GCC CTG GAG GGC GTC CCC CAG AAG AGC CGG 1240

```

EP 0 824 148 A2

	Leu	Leu	Ala	Ser	Val	Asp	Ala	Leu	Glu	Gly	Val	Pro	Gln	Lys	Ser	Arg	
				125					130					135			
5	GAG	CGG	CTG	CGT	TTC	TTC	ACC	CGC	CAG	TAC	GTC	AAC	GCC	ATG	GCC	CCC	1288
	Glu	Arg	Leu	Arg	Phe	Phe	Thr	Arg	Gln	Tyr	Val	Asn	Ala	Met	Ala	Pro	
				140					145					150			
	AGC	AAC	TTC	CTG	GCC	ACC	AAC	CCC	GAG	CTG	CTC	AAG	CTG	ACC	CTG	GAG	1336
	Ser	Asn	Phe	Leu	Ala	Thr	Asn	Pro	Glu	Leu	Leu	Lys	Leu	Thr	Leu	Glu	
				155					160					165			
	TCC	GAC	GCG	CAG	AAC	CTG	GTG	CGC	GGA	CTG	GCC	CTC	TTG	GCC	GAG	GAT	1384
10	Ser	Asp	Gly	Gln	Asn	Leu	Val	Arg	Gly	Leu	Ala	Leu	Leu	Ala	Glu	Asp	
						175					180				185		
	CTG	GAG	CGC	AGC	GCC	GAT	CAG	CTC	AAC	ATC	CGC	CTG	ACC	GAC	GAA	TCC	1432
	Leu	Glu	Arg	Ser	Ala	Asp	Gln	Leu	Asn	Ile	Arg	Leu	Thr	Asp	Glu	Ser	
						190					195				200		
	GCC	TTC	GAG	CTC	GGG	CGG	GAT	CTG	GCC	CTG	ACC	CCG	GCG	CGG	GTG	GTG	1480
15	Ala	Phe	Glu	Leu	Gly	Arg	Asp	Leu	Ala	Leu	Thr	Pro	Gly	Arg	Val	Val	
					205					210				215			
	CAG	CGC	ACC	GAG	CTC	TAT	GAG	CTC	ATT	CAG	TAC	AGC	CCG	ACT	ACC	GAG	1528
	Gln	Arg	Thr	Glu	Leu	Tyr	Glu	Leu	Ile	Gln	Tyr	Ser	Pro	Thr	Thr	Glu	
					220				225					230			
	ACG	GTG	GCG	AAG	ACA	CCT	GTG	CTG	ATA	GTG	CCG	CCC	TTC	ATC	AAC	AAG	1576
20	Thr	Val	Gly	Lys	Thr	Pro	Val	Leu	Ile	Val	Pro	Pro	Phe	Ile	Asn	Lys	
						240							245				
	TAC	TAC	ATC	ATG	GAC	ATG	CGG	CCC	CAG	AAC	TCC	CTG	GTG	GCC	TGG	CTG	1624
	Tyr	Tyr	Ile	Met	Asp	Met	Arg	Pro	Gln	Asn	Ser	Leu	Val	Ala	Trp	Leu	
						255					260				265		
	GTC	GCC	CAG	GCG	CAG	ACG	GTA	TTC	ATG	ATC	TCC	TGG	CGC	AAC	CCG	GCG	1672
	Val	Ala	Gln	Gly	Gln	Thr	Val	Phe	Met	Ile	Ser	Trp	Arg	Asn	Pro	Gly	
						270					275				280		
25	GTG	GCC	CAG	GCC	CAA	ATC	GAT	CTC	GAC	GAC	TAC	GTG	GTG	GAT	GCG	GTC	1720
	Val	Ala	Gln	Ala	Gln	Ile	Asp	Leu	Asp	Asp	Tyr	Val	Val	Asp	Gly	Val	
					285				290					295			
	ATC	GCC	GCC	GAC	GCG	GTG	GAG	GCG	GCC	ACC	GCG	GAG	GCG	GAG	GAG	GTG	1768
	Ile	Ala	Ala	Leu	Asp	Gly	Val	Glu	Ala	Ala	Thr	Gly	Glu	Arg	Glu	Val	
						300			305				310				
30	CAC	GCG	ATC	GCG	TAC	TGC	ATC	GCG	ACC	GCC	CTG	TCG	CTC	GCC	ATG		1816
	His	Gly	Ile	Gly	Tyr	Cys	Ile	Gly	Gly	Thr	Ala	Leu	Ser	Leu	Ala	Met	
						320					325						
	GCG	TGG	CTG	GCG	GCG	CGG	CAG	AAG	CAG	CGG	GTG	CGC	ACC	GCC	ACC		1864
	Gly	Trp	Leu	Ala	Ala	Arg	Arg	Gln	Lys	Gln	Arg	Val	Arg	Thr	Ala	Thr	
						335					340				345		
35	CTG	TTC	ACT	ACC	CTG	CTG	GAC	TTC	TCC	CAG	CCC	GGG	GAG	CTT	GCG	ATC	1912
	Leu	Phe	Thr	Thr	Leu	Leu	Asp	Phe	Ser	Gln	Pro	Gly	Glu	Leu	Gly	Ile	
						350					355				360		
	TTC	ATC	CAC	GAG	CCC	ATC	ATA	GCG	GCG	CTC	GAG	GCG	CAA	AAT	GAG	GCC	1960
	Phe	Ile	His	Glu	Pro	Ile	Ile	Ala	Ala	Leu	Glu	Ala	Gln	Asn	Glu	Ala	
					365				370					375			
40	AAG	GCG	ATC	ATG	GAC	GGG	CGC	CAG	CTG	GCG	GTC	TCC	TTC	AGC	CTG	CTG	2008
	Lys	Gly	Ile	Met	Asp	Gly	Arg	Gln	Leu	Ala	Val	Ser	Phe	Ser	Leu	Leu	
						380			385				390				
	CGG	GAG	AAC	AGC	CTC	TAC	TGG	AAC	TAC	TAC	ATC	GAC	AGC	TAC	CTC	AAG	2056
	Arg	Glu	Asn	Ser	Leu	Tyr	Trp	Asn	Tyr	Tyr	Ile	Asp	Ser	Tyr	Leu	Lys	
						395			400				405				
45	GGT	CAG	AGC	CCG	GTG	GCC	TTC	GAT	CTG	CTG	CAC	TGG	AAC	AGC	GAC	AGC	2104
	Gly	Gln	Ser	Pro	Val	Ala	Phe	Asp	Leu	Leu	His	Trp	Asn	Ser	Asp	Ser	
						415					420				425		
	ACC	AAT	GTG	GCG	GGC	AAG	ACC	CAC	AAC	AGC	CTG	CGC	CGT	CTC	TAC		2152
	Thr	Asn	Val	Ala	Gly	Lys	Thr	His	Asn	Ser	Leu	Leu	Arg	Arg	Leu	Tyr	
						430					435				440		
50	CTG	GAG	AAC	CAG	CTG	GTG	AAG	GGG	GAG	CTC	AAG	ATC	CGC	AAC	ACC	CGC	2200
	Leu	Glu	Asn	Gln	Leu	Val	Lys	Gly	Glu	Leu	Lys	Ile	Arg	Asn	Thr	Arg	
						445			450					455			
	ATC	GAT	CTC	GGC	AAG	GTG	AAG	ACC	CCT	GTG	CTG	CTG	GTG	TCG	GCG	GTG	2248
	Ile	Asp	Leu	Gly	Lys	Val	Lys	Thr	Pro	Val	Leu	Leu	Val	Ser	Ala	Val	
						460			465					470			

EP 0 824 148 A2

GAC GAT CAC ATC GCC CTC TGG CAG GGC ACC TGG CAG GGC ATG AAG CTG 2296
 Asp Asp His Ile Ala Leu Trp Gln Gly Thr Trp Gln Gly Met Lys Leu
 475 480 485
 TTT GGC GGG GAG CAG CGC TTC CTC CTG GCG GAG TCC GGC CAC ATC GCC 2344
 Phe Gly Gly Glu Gln Arg Phe Leu Leu Ala Glu Ser Gly His Ile Ala
 490 495 500 505
 GGC ATC ATC AAC CCG CCG GCC GCC AAC AAG TAC GGC TTT TGG CAC AAC 2392
 Gly Ile Ile Asn Pro Ala Ala Asn Lys Tyr Gly Phe Trp His Asn
 510 515 520
 GGG GCC GAG GCC GAG AGC CCG GAG AGC TGG CTG GCA GGG GCG ACC CAC 2440
 Gly Ala Glu Ala Glu Ser Pro Glu Ser Trp Leu Ala Gly Ala Thr His
 525 530 535
 CAG GGC GGC TCC TGG TGG CCC GAG ATG GGC TTT ATC CAG AAC CGT 2488
 Gln Gly Gly Ser Trp Trp Pro Glu Met Met Gly Phe Ile Gln Asn Arg
 540 545 550
 GAC GAA GGG TCA CAG CCC GTC CCC GCG CGG GTC CCG GAG GAA GGG CTG 2536
 Asp Glu Gly Ser Glu Pro Val Pro Ala Arg Val Pro Glu Glu Gly Leu
 555 560 565
 GCC CCC GCC CCC GGC CAC TAT GTC AAG GTG CCG CTC AAC CCC GTG TTT 2584
 Ala Pro Ala Pro Gly His Tyr Val Lys Val Arg Leu Asn Pro Val Phe
 570 575 580 585
 GCC TGC CCA ACA GAG GAG GAC GCC GCA TGAGCGCACA ATCCCTGGAA 2631
 Ala Cys Pro Thr Glu Glu Asp Ala Ala
 590
 GTAGGCCAGA AGGCCCGTCT CAGCAAGCGG TTCGGGGCGG CGGAGGTAGC CGCCTTCGCC 2691
 GCGCTCTCGG AGGACTTCAA CCCCTGCAC CTGGACCCGG CCTTCGCCGC CACCACGGCG 2751
 TTCGAGCGGC CCATAGTCCA CGGCATGCTG CTCGCCAGCC TCTTCTCCGG GCTGCTGGGC 2811
 CAGCAGTTGC CGGGCAAGGG GAGCATCTAT CTGGGTCAA GCCTCAGCTT CAAGCTGCCG 2871
 GTCTTTOTCG GGGACGAGGT GACGGCCGAG GTGGAGGTGA CCGCCCTTCG CGAGGACAAG 2931
 CCCATCGCCA CCTGACCAC CCGCATCTTC ACCCAAGGCG GCGCCCTCGC CGTGACGGGG 2991
 GAAGCCGTGG TCAAGCTGCC TTAAGCACCG GCGGCACGCA GGCACAATCA GCCCGGCCCC 3051
 TGCCGGGCTG ATTGTTCTCC CCGCTCCGC TTGCCCCCTT TTTCCGGGCA ATTTGGCCCA 3111
 GGCCCTTTCC CTGCCCGGCC TAACTGCCTA AAATGGCCGC CCTGCCGTGT AGGCATTCAT 3171
 CCAGCTAGAG GAATTC 3187

(2) INFORMATION FOR SEQ ID NO: 10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 3187 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 2611..3012

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

AGATCTGGAC CGGGGTGCTG GCCTGGGCCA CGCCGGCGAG GGCCAGCGCG GAGCAACCGA 60
 GCAGCAGGGC GAGAGGTTTC ATCGGGATTG CTGGCAGTC TGAATGACGT GCCAGCCTAT 120
 CAGCGCGGCG CCGGTGCGGC GAGGGCGCGC CGGACCCAGT CGGTACCTC TCGTCTGATC 180
 CGCCTCCCTC GACGGCGGTC GCTGACAAAA AAATTCAAAC AGAAATTAAC ATTTATGTCA 240
 TTTACACCAA ACCGCATTG GTTGACAGAT GCTCAAACGT GTGTTTGAAC AGAGCAAGCA 300
 ACACGTAAAC AGGGATGACA TGCAGTACCC GTAAGAAGGG CCGATTGGCC CACAACAACA 360
 CTGTTCTGCC GAACTGGAGA CCGATGATGA ATATGGACGT GATCAAGAGC TTTACCGAGC 420
 AGATGCAAGG CTTCGCGGCC CCCCTCACCC GCTACAACCA GCTGCTGGCC AGCAACATCG 480
 AACAGCTGAC CCGGTTGCAG CTGGCCTCCG CCAACGCCTA CGCCGAAGTG GGCCTCAACC 540
 AGTTGCAGGC CGTGAGCAAG GTGCAGGACA CCCAGAGCCT GGCGGCCCTG GGCACAGTGC 600
 AACTGGAGAC CGCCAGCCAG CTCTCCCGCC AGATGCTGGA TGACATCCAG AAGCTGAGCG 660

	CCCTCGGCCA	GCAGTTCAAG	GAAGAGCTGG	ATGTCTCTGAC	CGCAGACGGC	ATCAAGAAAA	720
	GCACGGGCAA	GGCCTGATAA	CCCTGGCTG	CCCGTTCCGG	CAGCCACATC	TCCCCATGAC	780
5	TCGACGCTAC	GGGCTAGTTC	CCGCCTCGGG	TGTGGGTGAA	GGAGAGCACA	TGAGCCAACC	840
	ATCTTATGGC	CCGCTGTTCC	AGGCCCTGGC	CCACTACAAT	GACAAGCTGC	TGGCCATGSC	900
	CAAGGCCAG	ACAGAGCGCA	CCGCCAGGC	GCTGCTGCAG	ACCAATCTGG	ACGATCTGGG	960
	CCAGGTGCTG	GAGCAGGGCA	GCCAGCAACC	CTGGCAGCTG	ATCCAGGCCC	AGATGAACTG	1020
	GTGGCAGGAT	CAGCTCAAGC	TGATGCAGCA	CACCTGTCTC	AAAAGGCGAG	GCCAGCCGAG	1080
	CGAGCCGGTG	ATCACCCCGG	AGCGCAGCGA	TCGCCGCTTC	AAGGCCGAGG	CCTGGAGCGA	1140
	ACAACCCATC	TATGACTACC	TCAAGCAGTC	CTACCTGTCT	ACCGCCAGGC	ACCTGCTGGC	1200
10	CTCGGTGGAT	GCCCTGGAGG	GCGTCCCCCA	GAAGAGCCGG	GAGCGGCTGC	GTTTCTTCAC	1260
	CCGCCAGTAC	GTCAACGCCA	TGGCCCCCAG	CAACTTCCTG	GCCACCAACC	CCGAGCTGCT	1320
	CAAGCTGACC	CTGGAGTCCG	ACGGCCAGAA	CCTGGTGGCG	GGACTGGCCC	TCTTGGCCGA	1380
	GGATCTGGAG	CGCAGCGCCG	ATCAGCTCAA	CATCCGCTTG	ACCGACGAAT	CCGCCTTCGA	1440
	GCTCGGGCGG	GATCTGGCCC	TGACCCCGGG	CCGGGTGGTG	CAGCGCACCG	AGCTCTATGA	1500
	GCTCATTCAG	TACAGCCCGA	CTACCGAGAC	GGTGGGCAAG	ACACCTGTGC	TGATAGTGCC	1560
15	GCCTTTCATC	AACAAGTACT	ACATCATGGA	CATGCGGCCC	CAGAACTCCC	TGGTCCGCTG	1620
	GCTGGTCCGC	CAGGGCCAGA	CGGTATTTCAT	GATCTCCTGC	CGCAACCCCG	GCGTGGCCCA	1680
	GGCCCAATC	GATCTCGAGC	ACTACGTGGT	GGATGGCGTC	ATCGCCGCCC	TGGACGGCGT	1740
	GGAGGCCGCC	ACCGCGCAGC	GGGAGGTGCA	CGGCATCGGC	TACTGCATCG	GCGGCACCGC	1800
	CCTGTCCGTC	GCCATGGGCT	GGCTGGCGGC	GCGGCGCCAG	AAGCAGCGGG	TGCGCACCGC	1860
	CACCCCTGTC	ACTACCTGTC	TGGACTTCTC	CCAGCCCGGG	GAGCTTGSCA	TCTTCATCCA	1920
	CGAGCCCATC	ATAGCGCGCG	TCGAGGCCGA	AAATGAGGCC	AAGGCGATCA	TGGACGGGCG	1980
20	CCAGCTGGCG	GTCTCCTTCA	GCCTGTCTGC	GGAGAACAGC	CTCTACTGGA	ACTACTACAT	2040
	CGACAGTTCG	CTCAAGGGTC	AGAGCCCGGT	GGCCTTCGAT	CTGCTGCACT	GGAACAGCGA	2100
	CAGCACCAAT	GTGGCGGGCA	AGACCCACAA	CAGCCTGCTG	CGCCGTCTCT	ACCTGGAGAA	2160
	CCAGCTGGTG	AAGGGGGAGC	TCAAGATCCG	CAACACCGGC	ATCGATCTCG	GCAAGGTGAA	2220
	GACCCCTGTG	CTGCTGGTGT	CGGCGGTGGA	CGATCACATC	GCCCTCTGGC	AGGGCACCTG	2280
	GCAGGGCATG	AAGCTGTTTG	GCGGGGAGCA	GCGCTTCCTC	CTGGCGGAGT	CCGGCCACAT	2340
25	CGCCGGCATC	ATCAACCCGC	CGGCCGCCAA	CAAGTACGGC	TTCTGGCACA	ACGGGGCCGA	2400
	GGCCGAGAGC	CCGGAGAGCT	GCGTGGCAGG	GGCGACGCAC	CAGGGCGGCT	CCTGCTGGCC	2460
	CGAGATGATG	GGCTTTATCC	AGAACCGTGA	CGAAGGGTCA	GAGCCCGTCC	CCGGCGGGGT	2520
	CCCGGAGGAA	GGGCTGGGCC	CCGCCCCCGG	CCACTATGTC	AAGGTGCGGC	TCAACCCCGT	2580
	GTTTGCCCTG	CCAACAGAGG	AGGACGCCGC	ATG AGC GCA CAA TCC CTG GAA GTA			2634
				Met Ser Ala Gln Ser Leu Glu Val			
30	GGC CAG AAG GCC CGT CTC AGC AAG CGG TTC GGG GCG GCG GAG GTA GCC						2682
	Gly Gln Lys Ala Arg Leu Ser Lys Arg Phe Gly Ala Ala Glu Val Ala						
	10 15 20						
	GCC TTC GCC GCG CTC TCG GAG GAC TTC AAC CCC CTG CAC CTG GAC CCG						2730
	Ala Phe Ala Ala Leu Ser Glu Asp Phe Asn Pro Leu His Leu Asp Pro						
	25 30 35 40						
35	GCC TTC GCC GCC ACC ACG GCG TTC GAG CGG CCC ATA GTC CAC GGC ATG						2778
	Ala Phe Ala Ala Thr Thr Ala Phe Glu Arg Pro Ile Val His Gly Met						
	45 50 55						
	CTG CTC GCC AGC CTC TTC TCC GGG CTG CTG GGC CAG CAG TTG CCG GGC						2826
	Leu Leu Ala Ser Leu Phe Ser Gly Leu Leu Gly Gln Gln Leu Pro Gly						
	60 65 70						
40	AAG GGG AGC ATC TAT CTG GGT CAA AGC CTC AGC TTC AAG CTG CCG GTC						2874
	Lys Gly Ser Ile Tyr Leu Gly Gln Ser Leu Ser Phe Lys Leu Pro Val						
	75 80 85						
	TTT GTC GGG GAC GAG GTG ACG GCC GAG GTG GAG GTG ACC GCC CTT CGC						2922
	Phe Val Gly Asp Glu Val Thr Ala Glu Val Glu Val Thr Ala Leu Arg						
	90 95 100						
45	GAG GAC AAG CCC ATC GCC ACC CTG ACC ACC CGC ATC TTC ACC CAA GGC						2970
	Glu Asp Lys Pro Ile Ala Thr Leu Thr Thr Arg Ile Phe Thr Gln Gly						
	105 110 115 120						
	GGC GCC CTC GCC GTG ACG GGG GAA GCC GTG GTC AAG CTG CCT						3012
	Gly Ala Leu Ala Val Thr Gly Glu Ala Val Val Lys Leu Pro						
	125 130						
50	TAAGCACCGG CGGCACGCAG GCACAATCAG CCCGCCCCCT GCCGGGCTCA TTGTTCTCCC						3072
	CCGCTCCGCT TGCCCCCTTT TTCGGGGCAA TTTGCCCCAG GCCCTTTCCC TGCCCCGCCT						3132
	AACTGCCTAA AATGGCCGCC CTGCCGTGTA GGCATTCTATC CAGCTAGAGG AATTC						3187

(2) INFORMATION FOR SEQ ID NO: 11:

EP 0 824 148 A2

- (1) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 25 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
 (A) DESCRIPTION: /desc = "synthetic DNA"

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

AGTTCCCGCC TCGGGTGTGG GTGAA

- (2) INFORMATION FOR SEQ ID NO: 12:

- (1) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 25 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
 (A) DESCRIPTION: /desc = "synthetic DNA"

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

GGCATATGCG CTCATGCCGC GTCCT 25

- (2) INFORMATION FOR SEQ ID NO: 13:

- (1) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 30 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
 (A) DESCRIPTION: /desc = "synthetic DNA"

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

GCCATATGAG CGCACAAATCC CTGGAAGTAG

- (2) INFORMATION FOR SEQ ID NO: 14:

- (1) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 30 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
 (A) DESCRIPTION: /desc = "synthetic DNA"

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

CTGGGATCCG CCGGTGCTTA AGGCAGCTTG

- (2) INFORMATION FOR SEQ ID NO: 15:

- (1) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

Ser Ala Gln Ser Leu Glu Val Gly Gln Lys Ala Arg Leu Ser Lys Arg
1 5 10 15
Phe Gly Ala Ala
20

(2) INFORMATION FOR SEQ ID NO: 16:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS:
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:

Met Ser Ala Gln Ser Leu Glu Val Gly Gln Lys Ala Arg Leu Ser Lys
1 5 10 15
Arg Phe Gly Ala Ala
20

Claims

1. A polyester synthase gene coding for a polypeptide containing the amino acid sequence of SEQ ID NO:2 or a sequence where in said amino acid sequence, one or more amino acids are deleted, replaced or added, said polypeptide bringing about polyester synthase activity.
2. A polyester synthase gene comprising the nucleotide sequence of SEQ ID NO:1.
3. A gene expression cassette comprising the polyester synthase gene of claims 1 or 2 and either of open reading frames located upstream and downstream of said gene.
4. The gene expression cassette according to claim 3, wherein the open reading frame located upstream of the polyester synthase gene comprises DNA coding for the amino acid sequence of SEQ ID NO:4.
5. The gene expression cassette according to claim 3, wherein the open reading frame located upstream of the polyester synthase gene comprises the nucleotide sequence of SEQ ID NO:3.
6. The gene expression cassette according to claim 3, wherein the open reading frame located downstream of the polyester synthase gene comprises DNA coding for a polypeptide containing the amino acid sequence of SEQ ID NO:6 or a sequence where in said amino acid sequence, one or more amino acids are deleted, replaced or added, said polypeptide bringing about enoyl-CoA hydratase activity.
7. The gene expression cassette according to claim 3, wherein the open reading frame located downstream of the


polyester synthase gene comprises the nucleotide sequence of SEQ ID NO:5.

8. A recombinant vector comprising the polyester synthase gene of claim 1 or 2 or the gene expression cassette of any one of claims 3 to 7.

9. A transformant transformed with the recombinant vector of claim 8.

10. A process for producing polyester, wherein the transformant of claim 9 is cultured in a medium and polyester is recovered from the resulting culture.

11. The process for producing polyester according to claim 10, wherein the polyester is a copolymer of 3-hydroxyalkanoic acid represented by formula I:



wherein R represents a hydrogen atom or a C1 to C4 alkyl group.

12. The process for producing polyester according to claim 10, wherein the polyester is a poly(3-hydroxybutyrate-co-3-hydroxyhexanoate) random copolymer.

FIG. 1A




FIG. 1B




FIG. 1C



FIG. 1D



FIG.2



Lane M: molecular-weight marker

Lane 1: soluble-protein fraction from NB3

Lane 2: active fraction eluted from the anion

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)